



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

darker ochraceous on flanks and sides of body; this difference of color is due partly, at least, to the worn condition of plumage of the latter.

One of the females of *C. robinsoni* differs from the other, and from the female of *C. phæniceus*, in having the scarlet vermilion of the crest, under part of wing and tail replaced by ochraceous yellow; otherwise it is the same.

The measurements of our pair of *C. phæniceus* (from Lake Maracaibo) are: ♂, wing, 3.45; tail, 3.60; tarsus, .94; culmen, .73; length of crest, 1.63 inches. ♀, wing, 3.33; tail, 3.53; tarsus, .97; culmen, .71; length of crest, 1.43 inches.

The specimens of *C. robinsoni* measure: ♂, wing, 3.26; tail, 3.29; tarsus, .96; culmen, .76; length of crest, 1.17 inches. ♂, wing, 3.20; tail, 3.20; tarsus, .95; culmen, .69; crest, 1.13 inches. ♂, wing, 3.20; tail, 3.10; tarsus, .95; culmen, .70; crest, 1.18 inches. ♀, wing, 3.13; tail, 3.06; tarsus, .91; culmen, .69; crest, 1.11 inches. ♀, wing, 3.20; tail, 3.13; tarsus, .99; culmen, .73; crest, 1.25 inches.

DESCRIPTIONS OF TWO NEW SUBSPECIES OF CALIFORNIA BIRDS.

BY F. STEPHENS.

Callipepla gambeli deserticola, subsp. nov. DESERT PARTRIDGE.

Subspecific characters.—Similar to *C. gambeli* but lighter in color; back olive gray inclining toward ash gray; sides chestnut, sharply striped with white; belly dull white; flanks and lower tail-coverts lightly streaked with brown; occiput russet in the male, drab in the female. Types, No. 4969, ♀ ad., 15 January, 1890, Palm Springs, San Diego Co. (now in Riverside Co.), Cal., alt. 100 ft.; and No. 4938, ♀ ad., 1 January, 1890, Walters, Colorado Desert, Cal., 250 feet below sea level. Both in my collection.

Habitat.—Desert District, from the western end of the Colorado Desert and the middle of the Mojave Desert east through the Colorado Valley.

The Desert Partridge differs from the Gambel's Partridge much as the Valley Partridge does from the California Partridge, and from the same cause, the drier habitat.

Speotyto cunicularia obscura, subsp. nov. DUSKY BUR-
ROWING OWL.

Subspecific characters.—Smaller than *S. c. hypogæa*; facial disk sepia, mixed with grayish; top of head sepia, streaked with dull brownish white; entire plumage slightly tinged with rusty. Length (of type), 8.65 inches (220 mm.); extent, 21.65 (550); wing, 6.20 (158); tail, 3.10 (79); tarsus, 1.65 (42). Type, No. 151,022, U. S. National Museum (original No. 5874), ♀ ad., 29 May, 1894, Upper Lake, Lake Co., Cal., alt. 1400 ft., F. Stephens, Col.

Habitat.—The valleys of the Pacific Coast District north of San Francisco Bay, perhaps excepting the lower valleys tributary to the Bay.

From the nature of the region Burrowing Owls are scarce all through it. The type is the only one I remember seeing. It is very dark about the head, especially about the face, which at first glance seems to be dirty, as if blood stained. The style of coloration is what might be expected in this region, being in the same direction as in *Bubo v. saturatus*, *Callipepla californica* and *Colaptes c. saturator*.